

Modelling the optically dark side of high-redshift galaxies

Bruno Guiderdoni¹

¹*Institut d'Astrophysique de Paris, CNRS, 98bis Boulevard Arago, F-75014 Paris.*

Abstract. The recent detection of the Cosmic Infrared Background in FIRAS and DIRBE residuals, and the observations of IR/submm sources by the ISOPHOT and SCUBA instruments have shed new light on the optically dark side of galaxy formation. It turns out that our view on the deep universe has been so far biased towards optically bright galaxies. We now know that a significant fraction of galaxy/star formation in the universe is hidden by dust shrouds. In this paper, we introduce a new modelling of galaxy formation and evolution that provides us with specific predictions in the IR/submm wavebands. These predictions are compared with the current status of the observations. Finally, the future all-sky and deep surveys with the PLANCK and FIRST missions are briefly described.

1 Introduction

The corrections needed to account for dust extinction in high-redshift galaxies are rather uncertain, and might induce an upward revision of the star formation rates (SFR) deduced from UV/visible observations by factors of a few ([10] [8]). Moreover, star formation might be completely hidden in heavily-extinguished galaxies which are missed at optical wavelengths. The discovery of the Cosmic Infrared Background in the FIRAS residuals between 200 μm and 2 mm ([11] [2], [1]) and in the DIRBE residuals at 240 and 140 μm ([4]) is a strong evidence that a significant fraction of star formation is hidden in dust shrouds. The level of the CIRB is more than five times higher than the “no-evolution” prediction based on the IRAS luminosity function, and twice as much as the level of the Cosmic Optical Background deduced from converging faint galaxy counts.

Recent observations at 850 μm with the SCUBA instrument of the JCMT ([14] [5]) have unveiled a large population of dusty objects. At the other end of the submm range, the FIRBACK program with ISOPHOT aboard ISO has yielded a catalogue of several hundred sources at 175 μm ([12]). The number counts are much higher than the “no-evolution” predictions. Preliminary attempts for identification seem to show that many of these objects are likely to be at $z > 1$.

We hereafter briefly describe a new modelling of galaxy formation ([2] [3]), and show predictions for on-going and forthcoming IR/submm surveys.




Figure 1: Predictions for faint galaxy counts at 15 μm , 60 μm , 175 μm , and 850 μm for a scenario fitting the CIRB, and data. 15 μm : ISO-HDF follow-up with ISOCAM ([9] solid squares), IRAS survey ([13] solid dot). 60 μm : IRAS survey ([7] open stars). 175 μm : ISOPHOT Lockman Hole ([6] open dot) and FIRBACK Marano field ([12] solid triangles). 850 μm : JCMT/SCUBA ([14] open squares).

2 Dust spectra in a semi-analytic framework

The IR/submm spectra of galaxies are computed in the following way: (1) follow chemical evolution of the gas; (2) implement extinction curves which depend on metallicity as in the Milky Way, the LMC and SMC; (3) compute τ_λ ; (4) assume the so-called “slab” geometry where the star and dust components are homogeneously mixed with equal height scales. (5) compute a spectral energy distribution by assuming a mix of various dust components. The contributions are fixed in order to reproduce the observational correlation of IRAS colours with total IR luminosity. These IR/submm spectra are implemented in a semi-analytic model of galaxy formation and evolution. This type of model has been very effective in computing the optical properties of galaxies in the paradigm of hierarchical clustering. We only extend this approach to the IR/submm range, and take the standard CDM case with $H_0=50$, $\Omega_0 = 1$, $\Lambda = 0$ and $\sigma_8 = 0.67$. We assume a Star Formation Rate $SFR(t) = M_{gas}/t_*$, with $t_* \equiv \beta t_{dyn}$. The efficiency parameter $1/\beta = 0.01$ gives a nice fit of local spirals. The robust result of this type of modelling is a rather flat cosmic star formation rate history. As a phenomenological way of reproducing the steep rise of the cosmic SFR history from $z = 0$ to $z = 1$, we introduce a “burst” mode of star formation with ten times higher efficiencies. This fairly




Figure 2: An $L_{bol} = 10^{12} L_{bol\odot}$ galaxy, similar to Arp 220, at various redshifts, compared to instrumental sensitivities for $t_{exp} = 1$ h.

reproduces the cosmic luminosity densities, as well as the CIRB.

Fig. 1 gives the predictions of number counts at 15, 60, 175 and 850 μm for the scenario which gives the best fit of the CIRB. The agreement of the predictions with the data seems good enough to suggest that these counts do probe the evolving population contributing to the CIRB. The model shows that 15 % and 60 % of the CIRB respectively at 175 μm and 850 μm are built up by objects brighter than the current limits of ISOPHOT and SCUBA deep fields. The predicted median redshift of the ISO-HDF is $z \sim 0.8$. It increases to $z \sim 1.5$ for the deep ISOPHOT surveys, and to $z \geq 2$ for SCUBA, even if the latter value seems to be very sensitive to the details of evolution.

3 Future surveys with PLANCK and FIRST

Fig. 2 gives a model spectral energy distribution typical of the $L_{bol} = 10^{12} L_{bol\odot}$ galaxy (similar to Arp 220) at various redshifts. The reader should note the specific behaviour of the observed flux at submm wavelengths, where the shift of the 60 – 100 μm rest-frame emission bump counterbalances distance dimming. The instrumental sensitivities of various on-going and forthcoming experiments are plotted on this diagram: the IRAS *Very Faint Source Survey*, ISOCAM, ISOPHOT, the IRAM interferometer, SCUBA, WIRE, SIRTf, SOFIA, the PLANCK *High Frequency Instrument*, the FIRST *Spectral and Photometric Imaging Receiver*, and the MMA/LSA.

The final sensitivity of these instruments is going to be confusion limited.

Table 1 gives the expected numbers of galaxies for the all-sky shallow survey of PLANCK *HFI*, and the medium-deep survey of FIRST *SPIRE* (to be launched by ESA in 2007), obtained from the model that fits the current counts. Faint counts produced by these two instruments are complementary. The study of the 250/350 and 350/500 colours are suited to point out sources which are likely to be at high redshifts. These sources can be eventually followed at 100 and 170 μm by the FIRST *Photoconductor Array Camera & Spectrometer* and by the FTS mode of *SPIRE*, to get the spectral energy distribution at $200 \leq \lambda \leq 600 \mu\text{m}$ with a typical resolution $R \equiv \lambda/\Delta\lambda = 20$.

Table 1. Predicted counts with PLANCK *HFI* and FIRST *SPIRE*.

λ	resolution	σ_{tot}	$N(> 5\sigma_{tot})$
<i>HFI</i> 350 μm	5'	180 mJy	5400*
<i>HFI</i> 550 μm	5'	100 mJy	3800*
<i>HFI</i> 850 μm	5'	50 mJy	3100*
<i>SPIRE</i> 250 μm	18"	3 mJy	70000**
<i>SPIRE</i> 350 μm	25"	3.2 mJy	60000**
<i>SPIRE</i> 500 μm	36"	4 mJy	27000**

*: 12 month nominal mission, 8 sr survey.

** : 3000 h medium-deep survey, 900 sec/pointing, 60 deg².

The FIRST satellite is ideally designed to probe the bulk of the 60 – 100 μm rest-frame emission of galaxies at $1 < z < 4$, and to recover the early SFR history of the universe. Our knowledge of the optically dark side of high-redshift galaxies should increase spectacularly in the forthcoming years.

References

- [1] Fixsen, D.J., *et al.*, 1998, ApJ , *in press*
- [2] Guiderdoni, B., *et al.*, 1997, Nature , 390, 257
- [3] Guiderdoni, B., *et al.*, 1998, MNRAS , 295, 877
- [4] Hauser, M.G., *et al.*, 1998, ApJ , *in press*
- [5] Hughes, D., *et al.*, 1998, Nature , *in press*
- [6] Kawara, K., *et al.*, 1998, A&A , *submitted*
- [7] Lonsdale, C.J., *et al.*, 1990, ApJ , 358, 60
- [8] Meurer, G.R., *et al.*, 1997, AJ , 114, 54
- [9] Oliver, S.J., *et al.*, 1997, MNRAS , 847, 1
- [10] Pettini, M., *et al.*, 1997, astro-ph/9707200
- [11] Puget, J.L., *et al.*, 1996, A&A , 308, L5
- [12] Puget, J.L., *et al.*, 1998, A&A , *submitted*
- [13] Rush, B., *et al.*, 1993, ApJS , 89, 1
- [14] Smail, I., *et al.*, 1997, ApJ , 490, L5